

18th International Symposium on Microbial Ecology (ISME), 14-19 August 2022

'Cold-adapted lipases and lipolytic bacteria found by metaproteogenomics in low-temperature anaerobic membrane bioreactors treating domestic wastewater'
 & **'Monitoring the anaerobic digestion microbial community as a foaming risk prediction method'**
 Authors: Reihaneh Bashiri *et al.*

I am the first author of two accepted abstracts (both as poster presentation) in the ISME 18th conference. One of the abstracts is from my PhD (*Cold-adapted lipases and lipolytic bacteria found by metaproteogenomics in low-temperature anaerobic membrane bioreactors treating domestic wastewater*) and the other is from the EBNet Proof of Concept (PoC) funding (*Monitoring the anaerobic digestion microbial community as a foaming risk prediction method*).

Both of these abstracts have value for the environmental or industrial biotechnology sector, and use combined omics tools: metagenomics to identify microbial individuals and metaproteomics to determine what are they doing within the communities of different environmental samples.

In my PhD research, I looked for the cold-adapted bacterial lipases and their producers in anaerobic treatment of domestic wastewater at low temperatures. Cold-adapted bacterial lipases have value for many industries like detergent manufacturers.

In the EBNet PoC, I looked for foam causing/stabilising bacteria and their biomarker molecules which correlates to foaming incidents in full-scale anaerobic digesters with a wide range of mixed solid and liquid wastes. This project can both offer a prediction tool for foaming incidents in anaerobic digestion systems and produce value for industries which are interested in biosurfactant (foam-causing substances with microbial origin) production by identifying novel biosurfactant producers.

Monitoring the anaerobic digestion microbial community as a foaming risk prediction method
 Reihaneh Bashiri¹, Lindsey Furness¹, Manpreet Bagga¹, Martin Pabst², Anil Wipat¹, Irina D. Ofiteru¹
¹School of Engineering, Newcastle upon Tyne, NE1 7RU, UK
²Deft University of Technology
³Newcastle University, School of Computing, Newcastle upon Tyne, NE2 4BW, UK
dana.ofiteru@newcastle.ac.uk

Foaming in anaerobic digestion (AD)

- Causes serious process instability
- Most foam-forming/stabilising bacteria in full-scale AD plant with mixed solid and liquid wastes are unknown

Methods

Hartlepool, Non-foaming plant
 Imperial Park, Foaming plant

Metagenomics
 MinION & Illumina
 Metaproteomics

Who are the putative foam-causing bacteria?
What are the putative foam-causing proteins?

Results

Who are the putative foam-forming bacteria and what are they doing? → Metaproteomics

Cold-adapted lipases and lipolytic bacteria found by metaproteogenomics
 in low-temperature anaerobic membrane bioreactors treating domestic wastewater
 Reihaneh Bashiri, Thomas P. Curtis, Irina D. Ofiteru
 Newcastle University, School of Engineering, Newcastle upon Tyne, NE1 7RU, UK
r.bashiri2@newcastle.ac.uk

Introduction

Cold-adapted bacteria can treat domestic wastewater at low temperatures. But poor lipid degradation is still a barrier.

Methods

We used metagenomics and metaproteomics [2].

Metagenomics → Who are the lipase producing bacteria?
 Metaproteomics → Are the lipases produced?
 Bioinformatics → KBase

Results

Who are the lipase producing bacteria?

- 40 lipolytic MAGs with 78 lipases
- Genome completeness ≥ 90%
- Contamination ≤ 10%
- At least one lipase (EC 3.1.1.3) gene

Out of 1519 total MAGs and 903 lipases (metagenome-assembled genomes)

How abundant are the lipolytic genera in reactors?

Non-stable 15°C, Stable 15°C, Non-stable 8°C, Stable 8°C

Who are the lipolytic genera from the metaproteomics side?

Metaproteomic found no lipase!
 Fatty (long-chain fatty acid transporter) was found as a lipolytic biomarker protein in 4 genera not recovered in lipolytic MAGs by metagenomics!

Conclusions

- Extracellular lipases and lipolytic bacteria were not easily identifiable by metagenomics and metaproteomics.
- Metaproteomics did not provide sufficient proteome coverage for lower abundant proteins such as lipases.
- Better protein extraction methods for metaproteomics of enzymes should be developed.

References

1. Pothof-Groen, E., et al. 2018. Environmental Science: Water Research & Technology, 4, 1002-1013.
 2. Bashiri, R. et al., 2021. Science Advances, 7(12), 1-15.